

GET THE FACTS ABOUT

MENSTRUATION

IN THIS ISSUE:

What is a "Normal" Period?
Tracking Your Period
Keepin' You Covered
PMS
Resources



WHAT IS A "NORMAL" PERIOD

No two people have the same period or cycle. As long as you aren't having any problems, you are most likely okay.

Below several examples of how periods can be different and still be normal.

men-strual-cy-cle:

The first time from the first day of one period to the first day of the next period.

- CYCLE LENGTHS can be from 21 to 45 days. Cycles can vary by a few days or be the same number of days every time.
- CYCLE REGULARITY fluctuates throughout your development. If you've just started menstruating, it's normal to have irregular cycles for the first year or so.
- DAYS OF BLEEDING can be as short as three days. Or five. Or anything in between.
- SYMPTOMS include cramps, back pain, or discomfort. More bowel movements than usual or constipation for the first day or so. There are also other symptoms (see PMS).

IS TOO MUCH BLOOD?

Sometimes there is worry about how much blood comes out. However, most days it only amounts to a teaspoon or two of blood. If you have any of the following problems, speak to your healthcare provider:


- blood that soaks through one or more pads or tampons every hour for several hours in a row.
- need to use pads and tampons at the same time to control the blood flow for several hours at a time.
- if you bleed or have your period more often than every 21 days



TRACKING YOUR PERIOD

BENEFITS OF TRACKING YOUR PERIOD

- You learn to understand your unique patterns
- You increase your awareness of your overall health
- You may develop a better understanding and management of moods



Did you know?

Your period is an indicator of your health, so knowing your pattern can help you know that everything is running fine. Tracking your period will help you remember details when speaking with your doctor.



pregnancy occurs

KEEPIN' YOU COVERED

What you choose to use to keep you covered during your period depends on what you are most comfortable using.



<u>PADS</u>

- Most commonly used
- Attaches to the inside of underwear the absorbent layer of cotton catches blood

TAMPONS

- Absorbs blood from inside vagina
- It is inserted into the vagina with an applicator or finger



MENSTRUAL CUPS

- Small, reusable, flexible cup that is inserted into the vagina and collects blood
- It may take a couple of tries before getting used to inserting it and removing it

MENSTRUAL PANTIES

- These are not regular panties. Menstrual panties are made with special fibers that absorb blood
- They are worn just like regular panties

PREMENSTRUAL SYNDROME-PMS

It is a term that refers to the physical and emotional symptoms one feels before or during their period

Typical symptoms include:

- Cramps
- Headaches
- Backaches
- Bloating or water retention
- Breast tenderness
- Pimples
- Mood swings including depression

Most symptoms are caused by hormonal changes with estrogen and progesterone levels. They tend to last a few days and are mild in nature. Most symptoms can be relieved by over-the-counter pain medication like acetaminophen (aka Tylenol).

FOR CRAMPS, TRY THIS:

- Light exercise
- A heating pad on the lowest setting
- Over-the-counter medications like acetaminophen or ibuprofen



MOST PMS SYMPTOMS ARE MILD AND LAST ONLY A FEW DAYS

RESOURCES

www.iknowmine.org
www.wernative.org
www.abovetheinfluence.org
www.projectknow.com
teens.drugabuse.gov
www.checkyourself.org

